

On a loss of information in a transition from quantum to a quasi-classical regime

Alex Granik*

Abstract

By defining information entropy in terms of probabilities densities $|\Psi|^2$ (Ψ is a wave function in the coordinate representation) it is explicitly shown how a loss of quantum information occurs in a transition from a quantum to a quasi-classical regime.

The Boltzmann-Gibbs-Shannon entropy

$$I = - \sum_{i=1}^N p_i \text{Log} p_i \quad (1)$$

(p_i is the classical probability) has an obvious relation to von Neumann entropy

$$S = -\text{Tr} \rho \text{Log} \rho$$

where the diagonal terms of the density matrix ρ are the probabilities p_i . Usually in quantum mechanics the probability p_i is considered as a function of energy E_i (e.g. [1]). Since we study a rather narrow problem of information dynamics when the probability is described in terms of the wave function Ψ , we will not discuss a more sophisticated (and complete) approach based on the density matrix and information processing involving quantum bits (qubits) (e.g., [2]).

To achieve our goal, we proceed along the lines used in derivation of continuity equation for probability density. To this end consider the probability in its

*Department of Physics, UOP, Stockton, CA. 95211; E-mail: agranik@uop.edu

coordinate representation. Thus p_i can be viewed as a probability of finding a particle in a spatial interval $\Delta q_i = q_{i+1} - q_i$:

$$p_i = \int_{q_i}^{q_{i+1}} \Psi(q)^* \Psi(q) dq \quad (2)$$

Using the mean value theorem, we rewrite Eq.(2) as follows:

$$p_i = |\Psi_i(\tilde{q}_i)|^2 \Delta q_i, \quad q_i \leq \tilde{q}_i \leq q_{i+1} \quad (3)$$

where the sum of p_i over all spatial intervals is

$$\sum_{i=1}^{\infty} p_i = \sum_{i=1}^{\infty} |\Psi_i(\tilde{q}_i)|^2 \Delta q_i = 1 \quad (4)$$

Using Eq.(3) we get

$$p_i \text{Log} p_i = |\Psi_i|^2 \Delta q_i [\text{Log}(|\Psi_i|^2) + \text{Log}(\Delta q_i)] \quad (5)$$

Now we take the following limit of (5)

$$\Delta q_i \rightarrow 0; \quad \Delta q_i \rightarrow dq$$

As a result, Eq.(5) yields

$$p_i \text{Log} p_i \rightarrow |\Psi(q)|^2 dq (\text{Log}|\Psi(q)|^2 - 1) + O(|\Delta q|^2) \quad (6)$$

and equation (1) becomes respectively:

$$I = - \int |\Psi(q)|^2 (\text{Log}|\Psi(q)|^2 - 1) dq \quad (7)$$

Therefore we narrow our discussion even more by restricting it with continuous spectra.

In a 3-D case the integration Eq.(7) is carried over a spatial volume d^3q . This means that the expression

$$\rho_I = -|\Psi(q)|^2 (\text{Log}|\Psi(q)|^2 - 1) \quad (8)$$

can be interpreted as the volume information density

$$\rho_I = dI/dV \quad (9)$$

We find its time variation:

$$\frac{\partial \rho_I}{\partial t} = -\frac{\partial |\Psi(q)|^2 (Log|\Psi(q)|^2 - 1)}{\partial t} = -\frac{\partial |\Psi|^2}{\partial t} Log|\Psi|^2 \quad (10)$$

If we use the continuity equation for probability density (e.g.,[3])

$$\frac{\partial |\Psi|^2}{\partial t} = -\frac{\hbar}{2im} div(|\Psi|^2 \nabla Log \frac{\Psi}{\Psi^*})$$

then we obtain from Eq.(10)

$$\begin{aligned} \frac{\partial \rho_I}{\partial t} &= \frac{\hbar}{2im} div(|\Psi|^2 \nabla Log \frac{\Psi}{\Psi^*}) Log|\Psi|^2 \equiv \\ &div(|\Psi|^2 Log|\Psi|^2 \frac{\hbar}{2im} \nabla Log \frac{\Psi}{\Psi^*}) - \frac{\hbar}{2im} \nabla Log \frac{\Psi}{\Psi^*} \bullet \nabla |\Psi|^2 \end{aligned} \quad (11)$$

Inserting the definition of the information density ρ_I , Eq.(8), and denoting the probability density $|\Psi|^2 = \rho$ into the first term on the right hand side of Eq.(11) , we obtain

$$\frac{\partial \rho_I}{\partial t} = -div[\frac{\hbar}{2im}(\rho_I - \rho) \nabla Log \frac{\Psi}{\Psi^*}] - \frac{\hbar}{2im} \nabla Log \frac{\Psi}{\Psi^*} \bullet \nabla |\Psi|^2 \quad (12)$$

Let us consider the transition to the classical case by representing the wave function as follows

$$\Psi = \sqrt{\rho} e^{iS/\hbar} \quad (13)$$

Using (13) in (12), we get

$$\frac{\partial \rho_I}{\partial t} = -div[\frac{\nabla S}{m}(\rho_I - \rho)] - \frac{\nabla S}{m} \bullet \nabla \rho \quad (14)$$

In the classical limit $\hbar t_c / m L_c^2 \rightarrow 0$ (where L_c is a characteristic length and t_c is the characteristic time) phase S becomes a classical action. As a result $\nabla S / m = \vec{v}$, that is a classical particle velocity. This means that in this limit Eq.(14) yields

$$\frac{\partial \rho_I}{\partial t} = -div[\vec{v}(\rho_I - \rho)] - \vec{v} \bullet \nabla \rho \quad (15)$$

If we integrate (15) over a volume V , use definition of the information entropy (9), and Gauss's theorem, we obtain

$$\frac{\partial}{\partial t} \int \rho_i dV = \frac{\partial I}{\partial t} = - \oint (\rho_I - \rho) \vec{v} \bullet d\vec{A} - \int \vec{v} \bullet \nabla \rho dV \quad (16)$$

By taking the boundary of the volume V to infinity and assuming that $\rho(q \rightarrow \infty) \rightarrow 0$ we obtain from (16)

$$\frac{\partial I}{\partial t} = - \int \vec{v} \bullet \nabla \rho dV \quad (17)$$

This means that in a transition from a quantum to classical regime the information entropy I [defined in the above narrow sense, Eq.(7)] is not conserved. Instead, if the classical velocity is in the direction of an increase of probability density, the information entropy decreases. Inversely, if the classical velocity is in the direction of a decreasing probability density, the information density increases.

These results are in agreement with the meaning of the information entropy as a number of states accessible to a system. In a transition to a classical regime this number drastically decreases, thus signaling a decrease in information entropy. Quite in agreement with that, and from another point of view, the less probable states carry more information than the more probable states. One can interpret these results as a statement that a classical regime has less degree of freedom than its quantum counterpart. This is definitely true when there is such a counterpart.

As we have already stated earlier, the above definition of information entropy is not fully appropriate for quantum mechanics because it does not take into account the information carried by a quantum phase. This serves as a strong indication that in the quantum region one needs to use another, more general, definition of information entropy, not necessarily associated with the qubits, which will account for the information associated with the quantum phase [4].

Acknowledgments

The author thanks V.Panico and C.Wulfman for the illuminating discussions of the results.

References

- [1] L.Landau and E.Lifshitz, Statistical Physics,Pergamon Press, 1980
- [2] N.Cerf and C.Adams, Negative Entropy in quantum Information Theory, quant-ph/9610005.
- [3] L.Landau and E.Lifshitz, Quantum Mechanics,Pergamon Press, 1974
- [4] A.Granik, G.Chapline, in preparation